



저작자표시-비영리-변경금지 2.0 대한민국

이용자는 아래의 조건을 따르는 경우에 한하여 자유롭게

- 이 저작물을 복제, 배포, 전송, 전시, 공연 및 방송할 수 있습니다.

다음과 같은 조건을 따라야 합니다:



저작자표시. 귀하는 원저작자를 표시하여야 합니다.



비영리. 귀하는 이 저작물을 영리 목적으로 이용할 수 없습니다.



변경금지. 귀하는 이 저작물을 개작, 변형 또는 가공할 수 없습니다.

- 귀하는, 이 저작물의 재이용이나 배포의 경우, 이 저작물에 적용된 이용허락조건을 명확하게 나타내어야 합니다.
- 저작권자로부터 별도의 허가를 받으면 이러한 조건들은 적용되지 않습니다.

저작권법에 따른 이용자의 권리는 위의 내용에 의하여 영향을 받지 않습니다.

이것은 [이용허락규약\(Legal Code\)](#)을 이해하기 쉽게 요약한 것입니다.

[Disclaimer](#)

Impact of pretreatment hydronephrosis on
the success rate of shock wave lithotripsy
in patients with ureteral calculi

Ki Don Chang

Department of Medicine

The Graduate School, Yonsei University

Impact of pretreatment hydronephrosis on the success rate of shock wave lithotripsy in patients with ureteral calculi

Directed by Professor Kang Su Cho

The Master's Thesis
submitted to the Department of Medicine,
the Graduate School of Yonsei University
in partial fulfillment of the requirements for the degree
of Master of Medical Science

Ki Don Chang

December 2016

This certifies that the Master's Thesis of
Ki Don Chang is approved.

Thesis Supervisor: Kang Su Cho

Thesis Committee Member#1: Joo Yong Lee

Thesis Committee Member#2: Sung Yoon Park

The Graduate School
Yonsei University

December 2016

ACKNOWLEDGEMENTS

For the past two years, my colleagues have continuously given me support and encouragement for this dissertation. I would like to express my deepest gratitude to all who have contributed to this work in various ways.

First of all, I would like to thank my thesis supervisor, Professor Kang Su Cho for being the most supportive mentor for me to finish my Master's thesis. His kind advices and patience have always been a great encouragement for me.

I also appreciate Professor Joo Young Lee who gave me a big hand and meaningful criticisms on this study. I probably couldn't make it without his teaching and tremendous encouragement. Likewise, I am also thankful to Professor Sung Yoon Park who gave me his expert advice to further enhance this manuscript. I would also like to thank all members of our department for their assistance and support.

Lastly, I thank my family from the bottom of my heart for their great support, love and being with me always.

<TABLE OF CONTENTS>

ABSTRACT	1
I. INTRODUCTION	3
II. MATERIALS AND METHODS	4
1. Patient cohort	
2. Good clinical practice protocols	
3. Shock wave lithotripsy	
4. Demographic data and stone characteristics	
5. On non-contrast computed tomography	
6. Hydronephrosis grading system	
III. RESULTS	7
IV. DISCUSSION	15
V. CONCLUSION	19
REFERENCES	21
ABSTRACT (IN KOREAN)	24

LIST OF FIGURES

Figure 1. Comparison of maximal stone length according to grade of hydronephrosis	11
Figure 2. Comparison of mean stone density according to grade of hydronephrosis	12

LIST OF TABLES

Table 1. Comparison of clinical characteristics between success and failure group after shock wave lithotripsy for ureteral stone	9
Table 2. Comparison of clinical characteristics among groups according to the grade of hydronephrosis	10
Table 3. Comparison of clinical characteristics between groups with pretreatment hydronephrosis grade 0-2 and grade 3-4	13
Table 4. Univariate and multivariate logistic regression of various clinical parameters for the prediction of one-session success after shock wave lithotripsy	14

ABSTRACT

Impact of pretreatment hydronephrosis on the success rate of shock wave lithotripsy in patients with ureteral calculi

Ki Don Chang

*Department of Medicine
The Graduate School, Yonsei University*

(Directed by Professor Kang Su Cho)

A retrospective study design was used to evaluate the predictors of the one-session success rate of shock wave lithotripsy (SWL). We also examined the relationships between pretreatment hydronephrosis grade and one-session SWL success rates. The medical records of 1,824 consecutive patients who underwent the first session of SWL for treatment of urinary stones between 2005 and 2013 were reviewed. After exclusion, 700 patients with single, 4–20 mm diameter radiopaque calculus were included in the study. The mean patient age was 52.55 ± 13.88 years. The mean maximal stone length (MSL) and skin-to-stone distance (SSD) were 9.24 ± 3.91 mm and 110.80 ± 18.98 cm, respectively. The average values for mean stone density (MSD) and stone heterogeneity index (SHI) were 707.04 ± 272.10 and 244.90 ± 110.16 , respectively. One-session success rate was 68.4%, 75.0%, 75.1%, 54.0%, and 10.5% in patient with hydronephrosis grade 0, 1, 2, 3, and 4, respectively. Patients were classified into success or failure groups based on SWL outcome. Multivariate logistic regression analyses revealed that MSL (odds ratio

(OR) 0.888, 95% confidence interval (CI): 0.841–0.934, $P<0.001$), MSD (OR 0.996, 95% CI: 0.995–0.997, $P<0.001$), SHI (OR 1.007, 95% CI: 1.005–1.010, $P<0.001$), and pretreatment hydronephrosis grade (OR 0.601, 95% CI: 0.368–0.988, $P=0.043$) were significantly associated with one-session success. Pretreatment grades 3 or 4 hydronephrosis were associated with failure of SWL in patients with single ureteral stone. In the presence of severe hydronephrosis, especially hydronephrosis grade 4, physicians should proceed cautiously in choosing and offering SWL as the primary treatment for ureteral stone.

Key words: shock wave lithotripsy, ureteral calculi, hydronephrosis

I. INTRODUCTION

Ureteral stone is one of the most common urologic diseases. Affected patients can experience extreme pain. There are many options for ureteral stone treatment (e.g., shock wave lithotripsy (SWL), and ureteroscopic, laparoscopic, or open surgery). Since the 1980s, SWL has been recommended as a non-invasive, effective, first-line treatment for small-sized radio-opaque stones (≤ 2 cm diameter).¹ The overall stone-free rate after treatment using SWL is 80–90%.^{2,3} Patient's age, sex, and stone characteristics (stone site, size, and density) are factors associated with stone-free rates. The number of stones, a history of urolithiasis, presence of renal colic, degree of hydronephrosis, and presence of a double J stent are additional factors associated with stone-free rates after treatment using SWL.⁴⁻⁶

Ureteral stone is the most common cause of ipsilateral hydronephrosis. The presences of one or more ureteral stones warrant urgent intervention to resolve the patient's symptoms and prevent damaging of renal function. The relationship between the degree of preoperative hydronephrosis and stone free rate after SWL has been studied. However, unanswered questions remain and a complete understanding would aid clinical decision-making during treatment selection.⁵⁻⁸ We aimed to investigate the predictors of one-session success rate after treatment using SWL; we specifically focused on the significance of pretreatment hydronephrosis in patients with a ureteral stone.

II. MATERIALS AND METHODS

1. Patient cohort

Medical records were obtained from a database of patients (N=1,824) who underwent an initial session of SWL between 1st November 2005 and 31st December 2014 at Severance Hospital, Seoul, Korea. The study inclusion criteria were: single, 4–20 mm, radiopaque calculus located within the ureter on plain-film X-rays, presentation within 1 month prior to SWL treatment, and no evidence of stone migration. Patients with bilateral ureteral stones, urinary tract congenital anomalies, or single kidney, and those who received prophylactic medical expulsion therapy, were excluded from the analysis. Data from a total of 700 patients were eligible for inclusion in the analysis.

2. Good clinical practice protocols

The study was performed in accordance with all applicable laws and regulations, good clinical practices, and the ethical principles described in the Declaration of Helsinki. The Institutional Review Board of Severance Hospital approved the study protocol (4-2016-0791). The study was exempt from the written informed consent requirement because it used a retrospective design and because the patients' records and information were anonymized and de-identified before analysis.

3. Shock wave lithotripsy

SWL was performed using an electro-conductive lithotripter (EDAP Sonolith Praktis, Technomed, and Lyon, France) until 31st December 2011. On 1st January 2012, this lithotripter was replaced by an electromagnetic generative lithotripter (Dornier Compact Delta II lithotripter, Dornier Medtech, Wessling, Germany). All SWL procedures were performed using fluoroscopic guidance. The total numbers of shocks ranged from 2500 to 4000 in each session, at a rate of 60–90 shock waves per minute, and a launch intensity ranging from 16 to 55 MPa.

4. Demographic data and non-contrast computed tomography stone characteristics

A detailed medical history that included the number of past stone events was obtained for each patient. Stone characteristics such as location, maximal stone length (MSL), stone heterogeneity index (SHI), skin-to-stone distance (SSD), and mean stone density (MSD) were evaluated. SSD was measured in the axial plane, 45° from the vertical axis.⁹ MSL was the longest stone length measured in three dimensions on non-contrast computed tomography (NCCT) images. We used the GE Centricity system (GE Healthcare Bio-Sciences Corp., Piscataway, NJ) for the measuring procedures. MSD was measured using bone windows on the magnified, axial NCCT image of the stone in the maximal diameter; the elliptical region of interest incorporated the largest cross-sectional area of the stone without including adjacent soft tissue.¹⁰ The SHI was defined as the standard deviation of the Hounsfield units (HU) in the same way used by Lee et al.¹¹ Successful SWL

treatment of the ureteral calculus was defined as the patient being rendered stone-free and asymptomatic, with clinically insignificant residual fragments ≤ 3 mm maximal diameter (measured by using a radiograph) within 2 weeks after single SWL treatment and not requiring additional treatment within a 3-month follow-up period.¹²

5. Hydronephrosis grading system

The hydronephrosis grading system was defined according to the degrees of change in the upper collecting system and used the Society of Fetal Ultrasound Grade system.¹³ Grade 0 hydronephrosis was defined as no dilatation of the renal pelvis, with calyceal walls opposed to each other. Grade 1 was defined as dilatation of the renal pelvis without dilatation of the calyces (could also be present in the extra renal pelvis), and no parenchymal atrophy. Grade 2 hydronephrosis was defined as dilatation of the renal pelvis (mild) and calyces (the pelvicalyceal pattern was retained), and no parenchymal atrophy. Grade 3 was defined as moderate dilatation of the renal pelvis and calyces, blunting of the fornices and flattening of the papillae, and possible mild cortical thinning. Grade 4 hydronephrosis was defined as gross dilatation of the renal pelvis and calyces with a ballooned appearance, loss of the borders between the renal pelvis, calyces, and renal atrophy indicated by the presence of cortical thinning.

6. Statistical analyses

The statistical comparisons of the continuous variables from the patients' demographic information were performed using the Student's or Welch's two-sample t-tests or the Wilcoxon rank sum test. One-way analysis of variance (ANOVA) was used subgroup analysis. After ANOVA, Tukey–Kramer's post hoc tests were used for between-group comparisons. Categorical variables were compared using Pearson's chi-square tests. Univariate and binomial multivariate logistic regression analyses were performed to define the factors that predicted post-SWL outcomes. The statistical analyses were performed using R software (version 3.0.3, R Foundation for Statistical Computing, Vienna, Austria; <http://www.r-project.org>).

III. RESULTS

Table 1 presents the results for the baseline characteristics of the 700 patients who underwent a first SWL treatment for single ureteral calculus. The mean patient age for the cohort was 52.55 ± 13.88 years. The mean MSL and SSD values were 9.24 ± 3.91 mm and 110.80 ± 18.98 cm, respectively. The mean MSD and SHI values were 707.04 ± 272.10 and 244.90 ± 110.16 HU, respectively. The results for pretreatment hydronephrosis grade in patients with ureteral calculi were 76 (10.9%) cases of grade 0, 383 (46.9%) cases of grade 1, 177 (25.3%) cases of grade 2, 100 (14.3%) cases of grade 3, and 19 (2.7%) cases of grade 4, hydronephrosis. There were 573 (81.9%) cases of upper ureteral stone, 48 (6.9%) cases of mid-ureteral

stone, and 79 (11.3%) cases of lower ureteral stone. The overall one-session success rate was 69.6%. The data from the study population were divided into two groups (treatment success or failure). The results indicated that there were statistical differences between the two groups in age (51.85 ± 13.88 years in success group versus 54.16 ± 13.80 years in failure group; $P=0.042$), MSL (8.38 ± 3.23 versus 11.23 ± 4.57 , respectively; $P<0.001$), MSD (642.63 ± 243.89 versus 854.30 ± 276.48 , respectively; $P<0.001$), SHI (252.06 ± 114.08 versus 228.51 ± 98.93 , respectively; $P=0.006$), and grade of hydronephrosis ($P<0.001$). The one-session success rates were 68.4%, 75.0%, 75.1%, 54.0%, and 10.5% for patients with hydronephrosis grades 0, 1, 2, 3, and 4, respectively (Table 2). The results indicated that there was a statistically significant association between hydronephrosis grade and the values for MSL and MSD ($P<0.001$; Table 2 and Figures 1 and 2). The associations between hydronephrosis grade and one-session success rate and stone location were also significantly different ($P<0.001$).

Table 1. Comparison of clinical characteristics between success and failure group after shock wave lithotripsy for ureteral stone

	Total cohort	Success group	Failure group	P-value
Patients, n	700	487	213	
Mean age \pm SD (yr)	52.55 \pm 13.88	51.85 \pm 13.88	54.16 \pm 13.80	0.042 ^a
Sex (%)				0.206 ^b
Male	454 (64.9)	308 (63.2)	146 (68.5)	
Female	246 (35.1)	179 (36.8)	67 (31.5)	
Mean MSL \pm SD (mm)	9.24 \pm 3.91	8.38 \pm 3.23	11.23 \pm 4.57	<0.001 ^a
Mean SSD \pm SD (cm)	110.80 \pm 18.98	110.66 \pm 18.90	11.15 \pm 19.21	0.751 ^a
Mean MSD \pm SD (HU)	707.04 \pm 272.10	642.63 \pm 243.89	854.30 \pm 276.48	<0.001 ^a
Mean SHI \pm SD (HU)	244.90 \pm 110.16	252.06 \pm 114.08	228.51 \pm 98.93	0.006 ^a
Hydronephrosis grade				<0.001
Grade 0	76 (10.9)	52 (10.7)	24 (11.3)	
Grade 1	328 (46.9)	246 (50.5)	82 (38.5)	
Grade 2	177 (25.3)	133 (27.3)	44 (20.7)	
Grade 3	100 (14.3)	54 (11.1)	46 (21.6)	
Grade 4	19 (2.7)	2 (0.4)	17 (8.0)	
Stone location, n (%)				0.510 ^b
Upper	573 (81.9)	403 (82.7)	170 (79.8)	
Middle	48 (6.9)	30 (6.2)	18 (8.5)	
Lower	79 (11.3)	54 (11.1)	25 (11.7)	
One-session success (%)	487 (69.6)	487 (69.6)	0 (0)	

^aBased on student's or Welch's two-sample t-tests ^bBased on Pearson's chi-squared tests with Yates' continuity correction FSF, first-time stone formers; HU, Hounsfield units; MSD, mean stone density; MSL, maximal stone length; n, number; RSF, recurrent stone formers; SD, standard deviation; SHI, stone heterogeneity index; SSD, skin-to-stone distance

Table 2. Comparison of clinical characteristics among groups according to the grade of hydronephrosis

	Grade 0	Grade 1	Grade 2	Grade 3	Grade 4	P-value
Patients, n	76	383	177	100	19	
Mean age \pm SD, (yr)	51.12 \pm 14.54	51.87 \pm 13.85	53.54 \pm 13.51	54.14 \pm 14.24	52.53 \pm 13.48	0.434
Sex (%)						0.029 ^b
Male	55 (72.4)	223 (68.0)	112 (63.3)	52 (52.0)	12 (63.2)	
Female	21 (27.6)	105 (32.0)	65 (36.7)	48 (48.0)	7 (36.8)	
Mean MSL \pm SD (mm)	7.99 \pm 2.80	8.49 \pm 3.55	9.29 \pm 3.52	11.74 \pm 4.82	13.65 \pm 3.32	<0.001 ^a
Mean SSD \pm SD (cm)	106.85 \pm 19.76	110.89 \pm 18.57	111.55 \pm 18.28	111.08 \pm 20.64	116.84 \pm 19.38	0.239 ^a
Mean MSD \pm SD (HU)	694.04 \pm 235.47	687.70 \pm 283.20	656.23 \pm 216.51	809.59 \pm 287.51	1026.54 \pm 273.84	<0.001 ^a
Mean SHI \pm SD (HU)	237.16 \pm 80.43	238.57 \pm 112.69	250.35 \pm 108.18	253.72 \pm 121.03	287.87 \pm 122.07	0.247 ^a
Stone location, n (%)						
Upper	56 (73.7)	283 (86.3)	145 (81.9)	73 (73.0)	16 (84.2)	
Middle	5 (6.6)	9 (2.7)	16 (9.0)	15 (15.0)	3 (15.8)	
Lower	15 (19.7)	36 (11.0)	16 (9.0)	12 (12.0)	0 (0.0)	
One-session success (%)	52 (68.4)	246 (75.0)	133 (75.1)	54 (54.0)	2 (10.5)	<0.001

^aBased on student's or Welch's two-sample t-tests ^bBased on Pearson's chi-squared tests with Yates' continuity correction FSF, first-time stone formers; HU, Hounsfield units; MSD, mean stone density; MSL, maximal stone length; n, number; RSF, recurrent stone formers; SD, standard deviation; SHI, stone heterogeneity index; SSD, skin-to-stone distance




Figure 1. Comparison of maximal stone length according to grade of hydronephrosis. It shows a statistically significant association between hydronephrosis grade and the values for MSL ($P < 0.001$).

The results also indicated that there were statistically significant differences in sex ($P=0.008$), MSL ($P<0.001$), MSD ($P<0.001$), stone location (<0.001), and one-session success rate ($P<0.001$) between the two groups with hydronephrosis grade 0–2 versus grades 3 and 4. The univariate logistic regression models revealed that the following predictive factors were associated with SWL outcome




Figure 2. Comparison of mean stone density according to grade of hydronephrosis. It shows a statistically significant association between hydronephrosis grade and the values for MSD ($P < 0.001$).

after ureteral stone treatment: age (OR: 0.988, 95% CI: 0.976–0.999; $P=0.043$), MSL (OR: 0.822, 95% CI: 0.782–0.861; $P<0.001$), MSD (OR: 0.997, 95% CI: 0.996–0.998; $P<0.001$), higher SHI (OR: 1.002, 95% CI: 1.001–1.004; $P=0.010$), and hydronephrosis grade (OR: 0.309, 95% CI: 0.206–0.463; $P<0.001$). The multivariate analyses revealed that shorter MSL, lower MSD, and higher SHI were independent predictors and hydronephrosis grades 3 and 4

were negative predictors of one-session success after SWL treatment for ureteral calculi (Table 4).

Table 3. Comparison of clinical characteristics between groups with pretreatment hydronephrosis grade 0-2 and grade 3-4

	Total cohort	Grade 0-2	Grade 3-4	P-value
Patients, n	700	581	119	
Mean age \pm SD (year)	52.55 \pm 13.88	52.28 \pm 13.84	53.88 \pm 14.07	0.251 ^a
Sex (%)				0.008 ^b
Male	454 (64.9)	390 (67.1%)	64 (53.8%)	
Female	246 (35.1)	191 (32.9%)	55 (46.2%)	
Mean MSL \pm SD (mm)	9.24 \pm 3.91	8.67 \pm 3.47	12.05 \pm 4.66	<0.001 ^a
Mean SSD \pm SD (cm)	110.80 \pm 18.98	110.56 \pm 18.67	112.00 \pm 20.47	0.451 ^a
Mean MSD \pm SD (HU)	707.04 \pm 272.10	678.94 \pm 258.54	844.23 \pm 295.23	<0.001 ^a
Mean SHI \pm SD (HU)	244.90 \pm 110.16	241.97 \pm 107.60	259.17 \pm 121.33	0.121 ^a
Stone location, n (%)				<0.001 ^b
Upper	573 (81.9)	484 (83.3%)	89 (74.8%)	
Middle	48 (6.9)	30 (5.2%)	18 (15.1%)	
Lower	79 (11.3)	67 (11.5%)	12 (10.1%)	
One-session success (%)	487	431 (74.2%)	56 (47.1%)	<0.001 ^b

^aBased on student's or Welch's two-sample t-tests

^bBased on Pearson's chi-squared tests with Yates' continuity correction

HU, Hounsfield units; MSD, mean stone density; MSL, maximal stone length; n, number; RSF, recurrent stone formers; SD, standard deviation; SHI, stone heterogeneity index; SSD, skin-to-stone distance

Table 4. Univariate and multivariate logistic regression of various clinical parameters for the prediction of one-session success aftershock wave lithotripsy

Parameter	OR	95% CI	P-value
Univariate			
Age	0.988	0.976–0.999	0.043
Sex (male)	0.789	0.558–1.110	0.177
MSL	0.822	0.782–0.861	<0.001
MSD	0.997	0.996–0.998	<0.001
SSD	0.999	0.990–1.007	0.750
SHI	1.002	1.001–1.004	0.010
Stone location			
Upper	Reference		
Middle	0.703	0.385–1.317	0.259
Lower	0.911	0.5b 54–1.533	0.719
Hydronephrosis Grade 3 & 4	0.309	0.206–0.463	<0.001
Multivariate			
Age	0.992	0.979–1.006	0.267
MSL	0.888	0.841–0.934	<0.001
MSD	0.996	0.995–0.997	<0.001
SHI	1.007	1.005–1.010	<0.001
Hydronephrosis Grade 3 & 4	0.601	0.368–0.988	0.043

HU, Hounsfield units; MSD, mean stone density; MSL, maximal stone length; n, number; RSF, recurrent stone formers; SD, standard deviation; SHI, stone heterogeneity index; SSD, skin-to-stone distance

IV. DISCUSSION

During the past decades, there have been significant changes in the management of urinary stone disease. Before the 1980s, open surgery treatment modality was used for most cases. This approach has been replaced by the use of minimally invasive surgical procedures such as ureterolithotomy, percutaneous nephrolithotomy, and SWL. Of these various options, SWL has been established as the preferred treatment for urinary stones due to its noninvasive character, few absolute contraindications, and resulting favorable clinical outcomes.^{14,15} However, if a satisfactory outcome for SWL is not expected for a specific clinical situation, then the other benefits (e.g., non-invasiveness) are no longer available to the patient. Therefore, it is essential to accurately predict individual treatment outcome in terms of proper treatment selection for patients who are candidates for SWL.

Previous studies of SWL success rates have revealed that age, sex, and stone characteristics (stone site, size, and density) are factors associated with post-treatment stone-free rates.^{5,8} Number of stones, a history of urolithiasis, the presence of renal colic, the degree of hydronephrosis, and use of a ureteral stent are other factors that affect success rate.⁶ MSL is a potentially useful independent predictor of SWL outcome.^{5,6} Patients with larger stones are more likely to experience failure of treatment and require further intervention.¹⁶ MSD is also an independent factor associated with SWL outcome.¹⁷ MSD is the mean

value of the HU of each pixel in a specific stone area and can be easily determined from the NCCT image using a picture archiving and communication system.¹⁸ This factor is also an independent predictor of SWL outcome.¹¹ SSD has been extensively investigated as a predictor of SWL success, but its use remains controversial for patients with ureter stones; SSD was a significant factor in one-half of all published studies.⁹ Recently, SHI was introduced as a novel independent predictor of SWL success in patients with ureteral stone and a useful clinical parameter for stone fragility.¹¹ Other studies of stone site found that treatment of proximal and distal ureteral stones results in better outcomes compared with treatment of mid-ureteral stones. However, the results of recent studies suggest that there is no differences between-group in stone site and treatment results.^{6,19} A history of urolithiasis has been reported as a negative factor affecting SWL success.^{20,21} Ureteral stenting is a significant factor affecting stone-free rates.¹⁹ Our study revealed that MSL, MSD, SHI, and severe hydronephrosis were significant predictors of one-session success rate after SWL treatment of single ureteral stone.

Several studies have evaluated the effects of preoperative hydronephrosis on the success rate of SWL; their findings have been inconsistent.⁵⁻⁸ El-Assmy et al. divided a total of 215 patients with single distal ureteral stone into two groups according to the absence or presence of hydronephrosis. There were no significant differences between the degree of stone-induced hydronephrosis and SWL outcome (83.2% in the non-hydronephrotic group versus 74.2% in

hydronephrotic group, $p=0.27$).⁷ They performed a similar study of 284 patients with proximal ureteral stone and found that the stone-free rate was 80.3% in the hydronephrotic group, compared with 89.1% in the patients without hydronephrosis ($p=0.12$).⁷ The results of Wang et al.'s multivariate analysis also suggested that hydronephrosis was not a significant factor for SWL treatment success rate (OR: 1.272, 95% CI: 0.471–3.433; $p=0.635$).⁶ In contrast, Kageyama et al. found that mid to lower ureteral calculi and moderate-to-severe hydronephrosis were negative predictive factors of SWL treatment success rate.⁸ Delakas et al. also found that the likelihood of SWL treatment failure increases with as the severity of the obstruction increases; hydronephrosis was associated with poorer results after SWL treatment (borderline significance; OR: 1.93, 95%CI: 0.99–3.77; $p=0.053$).⁵ The results of our multivariate analyses indicated that severe hydronephrosis (grades 3 & 4) was independent predictor of a poorer outcome after SWL treatment for ureteral calculi (Table 4). Generally, ureteral stones cause sudden ureteral obstruction that results in the development of hydronephrosis; continuous obstruction results in deterioration of renal function. Hydronephrosis has adverse effects on renal function and decreases ureteral peristalsis and hydraulic pressure. These changes may adversely affect the expulsion of ureteral stones. This relationship may help explain our results. Severe hydronephrosis can also be linked to ureteral stone impaction into the ureteral mucosa. Impacted stones are frequently associated with ureteral polyps or strictures. A chronically impacted stone may cause

inflammation and edema of the ureteral wall; these changes may also involve the surrounding tissues. The impacted stone can thus cause a more complete ureteral obstruction that result in a severe hydronephrosis. Impacted ureteral calculi are more difficult to fragment using SWL because of the lack of a natural expansion space for the targeted stones. There are currently two clinical definitions used for impacted stones.²² The first commonly used definition of impaction is the inability to pass a wire or catheter beyond the stone at the initial attempt.²³ The second definition of impaction is that the stone has remained at the same location in the ureter for more than 2 months.²⁴

Most urologists know that impacted stones are much more resistant to treatment using SWL. However, if currently available definitions for stone impaction are used, it is almost impossible to determine whether a specific ureteral stone is impacted at initial diagnosis. For these reasons, severe hydronephrosis can be used as ancillary clinical evidence of ureteral stone impaction in patients with ureteral stone when SWL is being considered as a primary treatment. European Association of Urology (EAU) guidelines recommend laparoscopic ureterolithotomy for treatment of large impacted stones when endoscopic lithotripsy or SWL has failed.²⁵ However, our study revealed that the one-session success rates were 54.0% and 10.5% in patient with hydronephrosis grades 3 and 4, respectively (Table 2). Therefore, even though SWL is less invasive compared with surgical lithotripsy; we are not sure whether physicians should offer SWL as a first-line therapy for ureteral stone patients with

concomitant grade 4 hydronephrosis. Physicians and patients should discuss this low SWL performance rate before selection of SWL as a treatment for ureteral stones with accompanying grade 4 hydronephrosis.

This study had some inherent limitations because the use of a retrospective study design may have introduced sampling bias. However, we used a relatively large cohort of patients who underwent SWL for treatment of single ureteral stone. The presence of renal stones and anatomical considerations including the location of calyx and renal pelvic stones or stones in the infundibulopelvic angle were additional possible sources of bias. To overcome this type of limitation and more clearly elucidate the effects of various stone-related factors on SWL outcomes, we limited the study population to patients who only had ureteral stones. Unlike previous studies, we also classified hydronephrosis grade into five groups using a current grading system. This approach resulted in a better characterization of the contribution of hydronephrosis to treatment outcome. Prospective studies that use large sample sizes are needed to confirm our results regarding the relationships between pretreatment hydronephrosis and stone clearance.

V. CONCLUSION

The presence of grades 3 or 4 hydronephrosis before SWL was a negative predictive factor for one-session success in patients treated for a single ureteral

stone. Severe hydronephrosis can be used as an indicator of possible ureteral stone impaction. In the presence of severe hydronephrosis, especially grade 4 hydronephrosis, physicians should be cautious when choosing and offering SWL as the primary treatment for ureteral stone.

REFERENCES

1. Frabboni R, Santi V, Ronchi M, Gaiani S, Costanza N, Ferrari G, et al. In situ echoguided extracorporeal shock wave lithotripsy of ureteric stones with the Dornier MPL 9000: a multicentric study group. *Br J Urol* 1994;73:487-93.
2. Creagh TA, Williams NN, Cronin K, Kerin MJ, Smith JM, Fitzpatrick JM. In situ ESWL for ureteric calculi: the optimum treatment? *Ir J Med Sci* 1993;162:348-50.
3. Singal RK, Denstedt JD. Contemporary management of ureteral stones. *Urol Clin North Am* 1997;24:59-70.
4. Srivastava A, Ahlawat R, Kumar A, Kapoor R, Bhandari M. Management of impacted upper ureteric calculi: results of lithotripsy and percutaneous litholapaxy. *Br J Urol* 1992;70:252-7.
5. Delakas D, Karyotis I, Daskalopoulos G, Lianos E, Mavromanolakis E. Independent predictors of failure of shockwave lithotripsy for ureteral stones employing a second-generation lithotripter. *J Endourol* 2003;17:201-5.
6. Wang M, Shi Q, Wang X, Yang K, Yang R. Prediction of outcome of extracorporeal shock wave lithotripsy in the management of ureteric calculi. *Urol Res* 2011;39:51-7.
7. El-Assmy A, El-Nahas AR, Youssef RF, El-Hefnawy AS, Sheir KZ. Impact of the degree of hydronephrosis on the efficacy of in situ extracorporeal shock-wave lithotripsy for proximal ureteral calculi. *Scand J Urol Nephrol* 2007;41:208-13.
8. Kageyama S, Hirai S, Higashi Y. [An investigation of factors associated with failure of extracorporeal shock wave lithotripsy for ureteral calculi]. *Hinyokika Kiyo* 2000;46:371-6.
9. Cho KS, Jung HD, Ham WS, Chung DY, Kang YJ, Jang WS, et al. Optimal Skin-to-Stone Distance Is a Positive Predictor for Successful Outcomes in Upper Ureter Calculi following Extracorporeal Shock Wave Lithotripsy: A Bayesian Model Averaging Approach. *PLoS One* 2015;10:e0144912.
10. Chung DY, Cho KS, Lee DH, Han JH, Kang DH, Jung HD, et al. Impact of colic pain as a significant factor for predicting the stone free rate of one-session shock wave lithotripsy for treating ureter stones: a Bayesian logistic regression model analysis. *PLoS One* 2015;10:e0123800.
11. Lee JY, Kim JH, Kang DH, Chung DY, Lee DH, Do Jung H, et al. Stone heterogeneity index as the standard deviation of Hounsfield units: A novel predictor for shock-wave lithotripsy outcomes in ureter calculi. *Sci Rep* 2016;6:23988.

12. Seitz C, Fajkovic H, Waldert M, Tanovic E, Remzi M, Kramer G, et al. Extracorporeal shock wave lithotripsy in the treatment of proximal ureteral stones: Does the presence and degree of hydronephrosis affect success? *Eur Urol* 2006;49:378-83.
13. Fernbach SK, Maizels M, Conway JJ. Ultrasound grading of hydronephrosis: introduction to the system used by the Society for Fetal Urology. *Pediatr Radiol* 1993;23:478-80.
14. Skolarikos A, Alivizatos G, de la Rosette J. Extracorporeal shock wave lithotripsy 25 years later: complications and their prevention. *Eur Urol* 2006;50:981-90; discussion 90.
15. Augustin H. Prediction of stone-free rate after ESWL. *Eur Urol* 2007;52:318-20.
16. Kim HH, Lee JH, Park MS, Lee SE, Kim SW. In situ extracorporeal shockwave lithotripsy for ureteral calculi: investigation of factors influencing stone fragmentation and appropriate number of sessions for changing treatment modality. *J Endourol* 1996;10:501-5.
17. Gupta NP, Ansari MS, Kesarvani P, Kapoor A, Mukhopadhyay S. Role of computed tomography with no contrast medium enhancement in predicting the outcome of extracorporeal shock wave lithotripsy for urinary calculi. *BJU Int* 2005;95:1285-8.
18. Lim KH, Jung JH, Kwon JH, Lee YS, Bae J, Cho MC, et al. Can stone density on plain radiography predict the outcome of extracorporeal shockwave lithotripsy for ureteral stones? *Korean J Urol* 2015;56:56-62.
19. Hofbauer J, Tuerk C, Hobarth K, Hasun R, Marberger M. ESWL in situ or ureteroscopy for ureteric stones? *World J Urol* 1993;11:54-8.
20. Kanao K, Nakashima J, Nakagawa K, Asakura H, Miyajima A, Oya M, et al. Preoperative nomograms for predicting stone-free rate after extracorporeal shock wave lithotripsy. *J Urol* 2006;176:1453-6; discussion 6-7.
21. Abe T, Akakura K, Kawaguchi M, Ueda T, Ichikawa T, Ito H, et al. Outcomes of shockwave lithotripsy for upper urinary-tract stones: a large-scale study at a single institution. *J Endourol* 2005;19:768-73.
22. Mugiya S, Ito T, Maruyama S, Hadano S, Nagae H. Endoscopic features of impacted ureteral stones. *J Urol* 2004;171:89-91.
23. Morgentaler A, Bridge SS, Dretler SP. Management of the impacted ureteral calculus. *J Urol* 1990;143:263-6.
24. Mugiya S, Nagata M, Un-No T, Takayama T, Suzuki K, Fujita K. Endoscopic management of impacted ureteral stones using a small caliber ureteroscope and a laser lithotripter. *J Urol* 2000;164:329-31.

25. Turk C, Petrik A, Sarica K, Seitz C, Skolarikos A, Straub M, et al. EAU Guidelines on Interventional Treatment for Urolithiasis. Eur Urol 2016;69:475-82.

요관 결석 환자에서 치료 전 수신증이 충격파쇄석술 성공율에 미치는 영향

<지도교수 조강수>

연세대학교 대학원 의학과

장기돈

지난 30년 동안 충격파쇄석술(shock wave lithotripsy; SWL)은 2cm 이하의 방사선비투과성 요관결석의 1차 치료로 권장되어 왔다. 본 연구는 요관 결석 환자에서 충격파 쇄석술의 성공율에 영향을 미치는 요인을 분석하고, 특히 치료 전 환자 수신증과 충격파쇄석술의 성공율간의 관계를 알아보고자 한다. 2005년부터 2013년까지 초회 충격파쇄석술을 시행 받은 환자 1,824명을 의무기록을 후향적으로 조사하였고, 최종적으로 4-20mm 크기의 방사선 비 투과성 단일 요관결석을 가진 700명의 환자 데이터를 분석하였다. 환자의 평균 나이는 52.5세였고 평균 maximum stone length (MSL)값과 skin-to-stone distance (SSD)값은 각각 9.24 ± 3.91 mm, 110.80 ± 13.88 cm였다. Mean stone density (MSD)와 stone heterogeneity (SHI)의 평균값은 각각 707.04 ± 272.10 Hounsfield unit 및 244.90 ± 110.16 였다. 수신증은 총 5등급으로 나누었으며 0등급이 76명 (10.9%), 1등급이 383명 (46.9%), 2등급이 177명 (25.3%), 3등급이 100명 (14.3%)이었고, 4등급은 19명 (2.7%)으로 나타났다. 전체 대상군을 충격파쇄석술 성공군과 실패군으로 나누어서 비교 분석하였다. 두 군의 나이 ($P=0.042$), MSL ($P<0.001$), MSD ($P<0.001$), SHI ($P=0.006$), 그리고 수신증 등급 ($P<0.001$)은 통계학적으로 차이를 보였다. 수신증 등급을 0-2등급 군과 3-4등급 군으로 나누어 비교하였을 때, MSL, MSD 및 결석의 위치는 유의한 차이를 보였고 ($P<0.001$), 충격파쇄석술 성공율에서도 의미있는 차이가 관찰되었다 ($P<0.001$). 다양한 임상인자들이 충격파쇄석술의 성공율에 미치는 영향을 알아보기 위한 다 변량 분석에서는 MSL (odds ratio (OR) 0.822, $P<0.001$), MSD (OR 0.996,

$P < 0.001$), SHI (OR 1.007, $P < 0.001$), 그리고 수신증 등급 (OR 0.601, $P = 0.043$)이 독립적인 예측인자임을 확인할 수 있었다. 결론적으로 3-4등급의 수신증을 동반한 요관결석에서는 충격파쇄석술 성공율이 감소함을 알 수 있었다. 따라서 심한 수신증을 동반한 요관결석 환자에서 충격파쇄석술을 선택함에 있어 신중할 기해야 한다.

핵심되는 말: 충격파쇄석술, 요관 결석, 수신증